
Oscillation Criteria for Higher Order Nonlinear Functional Difference Equations: Critical Study

S. Kaleeswari^{1*} and B. Selvaraj²

DOI: 10.9734/bpi/rsmcs/v1

Abstract

In this chapter some criteria for the oscillation of high order functional difference equation of the form

$$\Delta^2 \left(r(n) \left[\Delta^{(m-2)} y(n) \right]^\alpha \right) + q(n) f[y(g(n))] = 0,$$

where $\sum_{s=n_0}^{\infty} r^{-\frac{1}{\alpha}}(s) < \infty$ and $m > 1$

is discussed. Examples are given to illustrate the results.

Keywords: Functional difference equation; Nonlinear; non-oscillation; oscillation.

2010 Mathematics Subject Classification: 39A10.

1 Introduction

The notion of nonlinear difference equation was studied intensively by R. P. Agarwal [1] and oscillatory properties were discussed by R. P. Agarwal et al.[2, 3, 4]. Difference equations find a lot of applications in the natural sciences, technology and population dynamics [5, 6, 7].

Recently there has been a lot of interest in the study of oscillatory behavior of solutions of nonlinear difference equations. We can see this in [8-13]. Researchers carried out their researches on the oscillatory and asymptotic behavior of solutions of linear higher order difference equations. During the last two decades, many authors all over the world have taken keen interest in studying

oscillatory behavior of solutions of functional difference equations, due to its important applications in the field of science and computers. Such equations have extensive application in economics, physics, chemical technology, medicine, dynamic systems, optimal control and many other fields. So, we consider the high order nonlinear functional difference equation of the form

$$\Delta^2 \left(r(n) \left[\Delta^{(m-2)} y(n) \right]^\alpha \right) + q(n) f[y(g(n))] = 0, \quad (1.1)$$

¹Department of Mathematics, Nallamuthu Gounder Mahalingam College, Bharathiar University, Coimbatore, Tamil Nadu, India.

²Department of Science and Humanities, Nehru Institute of Engineering and Technology, T. M. Palayam, Coimbatore, Tamil Nadu, India.

*Corresponding author: E-mail: kaleesdesika@gmail.com;

where

$$\sum_{s=n_0}^{\infty} r^{-\frac{1}{\alpha}}(s) < \infty. \quad (1.2)$$

subject to the hypotheses:

- (i) α is the ratio of any two positive odd integers.
- (ii) $\{r(n)\}, \{q(n)\}$ are real-valued positive sequences.
- (iii) $\{g(n)\}$ is a realvalued increasing sequence with $g(n) < n$, for $n \geq n_0$ and $\lim_{n \rightarrow \infty} g(n) = \infty$.
- (iv) $f: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function such that $xf(x) > 0$ and $f'(x) \geq 0$, for $x \neq 0$ and

$$-f(-xy) \geq f(xy) \geq f(x)f(y), \text{ for } xy > 0. \quad (1.3)$$

Here Δ is the forward difference operator defined by

$$\Delta y(n) = y(n+1) - y(n).$$

By a solution of equation (1.1), we mean a real sequence $\{y(n)\}$ which is defined for $n \geq \min_{i \geq 0} \{g(i)\}$ and satisfies equation (1.1) for all large n . A solution $\{y(n)\}$ is said to be oscillatory if it is neither eventually positive nor eventually negative. Otherwise it is called non-oscillatory. A difference equation is said to be oscillatory if all of its solutions are oscillatory. Otherwise, it is non-oscillatory.

Our aim in this chapter is to obtain some oscillation criteria for the solution of equation (1.1). For recent contributions to this study, we refer the reader to the papers [8-13] and the references cited therein.

2 Main Results

In this section, we are interested to present some sufficient conditions for the oscillation of all solutions of the equation (1.1).

Assume that conditions (i)-(iv), (1.2) and (1.3) hold. If the first order delay difference equation

$$\Delta u(n) + cq(n)f[(g(n) - n_2)^{m-3}]f\left[\sum_{s=n_2}^{g(n)-1} \left(\frac{s}{r(s)}\right)^{\frac{1}{\alpha}}\right]f\left[\left(u^{\frac{1}{\alpha}}(g(n))\right)\right] = 0, \quad (2.1)$$

for any constant $c, 0 < c < 1$ and for all $N \geq n_0$ is oscillatory, then the equation (1.1) is oscillatory.

Proof. Let $\{y(n)\}$ be a non-oscillatory solution of equation (1.1). Without loss of generality, assume that $y(n) > 0$ and $y(g(n)) > 0$, for $n \geq n_0 \geq 0$.

Then from (1.1), we can see that

$$\Delta^2 \left(r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right) \leq 0, \text{ for } n \geq n_0. \quad (2.2)$$

There exists a $n_1 \geq n_0$ such that $\left\{ \Delta^{(m-2)} y(n) \right\}$ and $\left\{ \Delta \left(r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right) \right\}$ are eventually monotone and one-signed, for $n \geq n_1$.

We consider the following four cases for $n \geq n_1$ and $m > 1$.

- (i) $\Delta \left(r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right) > 0$ and $\Delta^{(m-2)} y(n) < 0$, for $n \geq n_1$.
- (ii) $\Delta \left(r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right) > 0$ and $\Delta^{(m-2)} y(n) > 0$, for $n \geq n_1$.
- (iii) $\Delta \left(r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right) < 0$ and $\Delta^{(m-2)} y(n) > 0$, for $n \geq n_1$.
- (iv) $\Delta \left(r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right) < 0$ and $\Delta^{(m-2)} y(n) < 0$, for $n \geq n_1$.

Case (i): Since $r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha$ is increasing, for $n \geq s \geq n_1$, we obtain

$$r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \geq r(s) \left(\Delta^{(m-2)} y(s) \right)^\alpha.$$

That is,

$$r(s) \left(\Delta^{(m-2)} y(s) \right)^\alpha \leq r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha,$$

which implies,

$$\Delta^{(m-2)} y(s) \leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}.$$

Summing the above inequality from n_1 to $n - 1$, we get

$$\Delta^{(m-3)} y(n) - \Delta^{(m-3)} y(n_1) \leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}.$$

That is,

$$\Delta^{(m-3)} y(n) \leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}} + \Delta^{(m-3)} y(n_1).$$

Since $\Delta^{(m-2)} y(n) < 0$, we have

$$\begin{aligned} \Delta^{(m-3)} y(n) &\leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}} \\ &= \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1) \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}. \end{aligned}$$

That is,

$$\Delta^{(m-3)} y(n) \leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1) \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}.$$

Similarly, we can obtain

$$\Delta^{(m-4)} y(n) \leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1)^2 \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}.$$

Proceeding like this, we get

$$\Delta y(n) \leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1)^{m-3} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}$$

and

$$y(n) \leq \left[r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1)^{m-3} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}. \quad (2.3)$$

From (1.2) and since $\Delta^{(m-2)} y(n) < 0$, we can see that $y(n) \rightarrow -\infty$ as $n \rightarrow \infty$, which contradicts our assumption that $y(n) > 0$.

Case (ii):

Let

$$z(n) = r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha.$$

Then $z(n) > 0$. Since $\Delta z(n)$ is decreasing (from (2.2)), for $n \geq n_1$, we have

$$z(n) - z(n_1) = \sum_{s=n_1}^{n-1} \Delta z(s) \geq (n - n_1) \Delta z(n).$$

Then there exists a $n_2 \geq n_1$ and a constant $b, 0 < b < 1$ such that

$$z(n) \geq bn \Delta z(n), \text{ for } n \geq n_2,$$

which implies

$$r(n) \left(\Delta^{(m-2)} y(n) \right)^\alpha \geq bn \Delta z(n).$$

That is,

$$\Delta^{(m-2)} y(n) \geq b^{\frac{1}{\alpha}} \left(\frac{n}{r(n)} \right)^{\frac{1}{\alpha}} (\Delta z(n))^{\frac{1}{\alpha}}, \text{ for } n \geq n_2.$$

Summing the above inequality from n_2 to $n - 1$, we get

$$\Delta^{(m-3)} y(n) \geq b^{\frac{1}{\alpha}} (n - n_2) \left(\frac{s}{r(s)} \right)^{\frac{1}{\alpha}} (\Delta z(n))^{\frac{1}{\alpha}}.$$

Similarly, we can obtain

$$\Delta y(n) \geq b^{\frac{1}{\alpha}} (n - n_2)^{m-3} \left(\frac{s}{r(s)} \right)^{\frac{1}{\alpha}} (\Delta z(n))^{\frac{1}{\alpha}}.$$

Also,

$$y(n) \geq b^{\frac{1}{\alpha}} (n - n_2)^{m-3} \sum_{s=n_2}^{n-1} \left(\frac{s}{r(s)} \right)^{\frac{1}{\alpha}} (\Delta z(n))^{\frac{1}{\alpha}}.$$

There exists a $n_3 \geq n_2$ such that

$$y(g(n)) \geq b^{\frac{1}{\alpha}} (g(n) - n_2)^{m-3} \sum_{s=n_2}^{g(n)-1} \left(\frac{s}{r(s)} \right)^{\frac{1}{\alpha}} (\Delta z(g(n)))^{\frac{1}{\alpha}}, \text{ for } n \geq n_3. \quad (2.4)$$

Using (2.4) and (1.3) in equation (1.1), we obtain

$$\begin{aligned} -\Delta^2 z(n) &= q(n) f(y(g(n))) \\ &\geq f\left[b^{\frac{1}{\alpha}}\right] q(n) f[(g(n) - n_2)^{m-3}] \\ &\quad f\left[\sum_{s=n_2}^{g(n)-1} \left(\frac{s}{r(s)}\right)^{\frac{1}{\alpha}}\right] f\left[(\Delta z(g(n)))^{\frac{1}{\alpha}}\right], \text{ for } n \geq n_3. \end{aligned}$$

Setting $u(n) = \Delta z(n) > 0$, for $n \geq n_3$ in the above inequality, we get

$$\begin{aligned} -\Delta u(n) &\geq f\left[b^{\frac{1}{\alpha}}\right] q(n) f[(g(n) - n_2)^{m-3}] \\ &\quad f\left[\sum_{s=n_2}^{g(n)-1} \left(\frac{s}{r(s)}\right)^{\frac{1}{\alpha}}\right] f\left[\left(u^{\frac{1}{\alpha}}(g(n))\right)\right]. \end{aligned}$$

That is,

$$\Delta u(n) + cq(n) f[(g(n) - n_2)^{m-3}] f\left[\sum_{s=n_2}^{g(n)-1} \left(\frac{s}{r(s)}\right)^{\frac{1}{\alpha}}\right] f\left[\left(u^{\frac{1}{\alpha}}(g(n))\right)\right] \leq 0, \quad (2.5)$$

where $c = f\left(b^{\frac{1}{\alpha}}\right)$, for $n \geq n_3$. Now the inequality (2.5) has eventually positive solution $u(n)$.

By a well-known result (see [10], [13]), the difference equation (2.1) also has an eventually positive solution which contradicts our assumption that (2.1) is oscillatory.

Case (iii):

This case cannot hold.

Let

$$z(n) = r(n) \left(\Delta^{(m-2)}y(n)\right)^\alpha.$$

Then $\Delta z(n) < 0$ and from (2.2), $\Delta^2 z(n) < 0$.

Therefore

$$\lim_{n \rightarrow \infty} z(n) = -\infty,$$

which is a contradiction to the fact that $\Delta^{(m-2)}y(n) > 0$.

Case (iv):

Since $r(n) \left(\Delta^{(m-2)}y(n)\right)^\alpha$ is decreasing, for $s \geq n \geq n_1$, we get

$$r(s) \left(\Delta^{(m-2)}y(s)\right)^\alpha \leq r(n) \left(\Delta^{(m-2)}y(n)\right)^\alpha,$$

which implies,

$$\Delta^{(m-2)}y(s) \leq \left[r(n) \left(\Delta^{(m-2)}y(n)\right)^\alpha\right]^{\frac{1}{\alpha}} \left(\frac{1}{r(s)}\right)^{\frac{1}{\alpha}}.$$

Summing the above inequality from n_1 to $n - 1$, we get

$$\Delta^{(m-3)}y(n) - \Delta^{(m-3)}y(n_1) \leq \left[r(n) \left(\Delta^{(m-2)}y(n)\right)^\alpha\right]^{\frac{1}{\alpha}} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)}\right)^{\frac{1}{\alpha}}.$$

That is,

$$\Delta^{(m-3)}y(n) \leq \left[r(n) \left(\Delta^{(m-2)}y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}} + \Delta^{(m-3)}y(n_1).$$

Since $\Delta^{(m-2)}y(n) < 0$, we have

$$\begin{aligned} \Delta^{(m-3)}y(n) &\leq \left[r(n) \left(\Delta^{(m-2)}y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}} \\ &= \left[r(n) \left(\Delta^{(m-2)}y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1) \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}. \end{aligned}$$

That is,

$$\Delta^{(m-3)}y(n) \leq \left[r(n) \left(\Delta^{(m-2)}y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1) \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}.$$

Similarly, we can obtain

$$\Delta^{(m-4)}y(n) \leq \left[r(n) \left(\Delta^{(m-2)}y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1)^2 \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}.$$

Proceeding like this, we get

$$\Delta y(n) \leq \left[r(n) \left(\Delta^{(m-2)}y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1)^{m-3} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}$$

and

$$y(n) \leq \left[r(n) \left(\Delta^{(m-2)}y(n) \right)^\alpha \right]^{\frac{1}{\alpha}} (n - n_1)^{m-3} \sum_{s=n_1}^{n-1} \left(\frac{1}{r(s)} \right)^{\frac{1}{\alpha}}. \quad (2.6)$$

From (1.2) and since $\Delta^{(m-2)}y(n) < 0$, we can see that $y(n) \rightarrow -\infty$ as $n \rightarrow \infty$, which contradicts our assumption that $y(n) > 0$. This completes the proof. \square

The following examples are illustrative.

3 Example

Example 3.1. Consider the fourth order equation

$$\Delta^2 \left(n^6 \left(\Delta^2 y(n) \right)^3 \right) + 64 \left[(n+2)^6 - n^6 \right] y^3(n-2) = 0. \quad (3.1)$$

Here $r(n) = n^6$, $\alpha = 3$ and $m = 4$.

Also,

$$\sum_{s=n_0}^{n-1} r^{-\frac{1}{\alpha}}(s) = \sum_{s=n_0}^{n-1} (n^6)^{-\frac{1}{3}} = \sum_{s=n_0}^{n-1} n^{-2} = \sum_{s=n_0}^{n-1} \frac{1}{n^2} < \infty$$

and $g(n) = n - 2 < n$.

We can easily see that all conditions of Theorem 2.1 are satisfied and hence all the solutions of equation (3.1) are oscillatory.

One of such solution is $y(n) = (-1)^n$.

Example 3.2. Consider the third order equation

$$\Delta^2 (n^9 (\Delta y(n))^3) + 216 [729 (n+2)^9 + 54 (n+1)^9 + n^9] y^3 (n-2) = 0. \quad (3.2)$$

Here $r(n) = n^9$, $\alpha = 3$ and $m = 3$.

Also,

$$\sum_{s=n_0}^{n-1} r^{-\frac{1}{\alpha}}(s) = \sum_{s=n_0}^{n-1} (n^9)^{-\frac{1}{3}} = \sum_{s=n_0}^{n-1} n^{-3} = \sum_{s=n_0}^{n-1} \frac{1}{n^3} < \infty$$

and $g(n) = n - 2 < n$.

We can easily see that all conditions of Theorem 2.1 are satisfied and hence all the solutions of equation (3.2) are oscillatory.

One of the solutions is $y(n) = \frac{(-3)^n}{2}$.

4 Conclusion

In this chapter, we have proposed the comparison method for identifying oscillatory solutions of higher order functional difference equations. This method compares first order equation which is very simple. Moreover, the above examples reveal the efficiency of our method.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Agarwal RP. Difference equations and inequalities Theory, Methods and Applications, 2nd edition, New York; 2000.
- [2] Agarwal RP, Martin Bohner, Said R. Grace, Donal O'Regan. Discrete oscillation theory. CMIA Book Series. ISBN: 977-5945-19-4.
- [3] Agarwal RP, Said R. Grace, Donal O'Regan. Oscillation theory for difference and functional differential equations. Kluwer Academic Publishers, Dordrecht; 2000.
- [4] Agarwal RP, Wong PJY. Advanced topics in difference equations. Kluwer Academic Publishers, Dordrecht; 1997.
- [5] Gyori I, Ladas G. Oscillation theory of delay differential equations with applications. Clarendon Press, Oxford; 1991.
- [6] Kelley WG, Peterson AC. Difference Equations an introduction with applications. Academic Press, Boston; 1991.
- [7] Lakshmikantham V, Trigiante D. Theory of difference equations, numerical methods and applications. Academic Press, New York; 1988.
- [8] Bolat Y, Akin O. Oscillatory behaviour of a higher order nonlinear neutral type functional difference equation with oscillating coefficients. Applied Mathematics Letters. 2004;17:1073-1078.
- [9] Grzegorzczuk G, Werbowski J. Oscillation of higher order linear difference equations. Computers and Mathematics with Applications. 2001;42:711-717.

- [10] Ladas G, Ch G. Philos, Sficas YG. Sharp conditions for the oscillation of delay difference equations. *Journal of Applied Mathematical Simulation*. 1989;2:101-111.
- [11] Selvaraj B, Kaleeswari S. Oscillation theorems for certain fourth order non-linear difference equations. *International Journal of Mathematics Research*. 2013;5(3):299-312.
- [12] Thandapani E, Selvaraj B. Existence and asymptotic behavior of non oscillatory solutions of certain nonlinear difference equations. *Far East Journal of Mathematical Sciences (FJMS)*. 2004;14(1):9-25.
- [13] Zhang BG, Yong Zhou. Oscillation of difference equations with several delays. *Computers and Mathematics with Applications*. 2002;44:817-821.

Biography of author(s)



S. Kaleeswari

Department of Mathematics, Nallamuthu Gounder Mahalingam College, Bharathiar University, Coimbatore, Tamil Nadu, India.

She is an Assistant Professor in the Department of Mathematics, Nallamuthu Gounder Mahalingam College, Pollachi. She has 14 years of experience in teaching and doing her research in the field of Difference and Differential Equations. She has published 13 papers in national and international journals. She published her results on oscillation theory of ordinary and delay Difference Equations. She has life membership in Indian mathematical society, Ramanujan mathematical society, Indian science congress association. She acts as reviewer for some refereed journals.



B. Selvaraj

Department of Science and Humanities, Nehru Institute of Engineering and Technology, T. M. Palayam, Coimbatore, Tamil Nadu, India.

He is Dean of Nehru Institute of Engineering and Technology, T. M. Palayam, Coimbatore. He has 44 years of experience in teaching and doing his research in the field of Difference and Differential Equations. He has published 55 papers in national and international journals and 4 books. He

published her results on oscillation theory of Difference Equations. He is a member of Editorial Board in Malaya Journal of Matematik, Bulletin of Pure & Applied Science and IOSR Journal.

© Copyright (2020): Authors. The licensee is the publisher (Book Publisher International).

DISCLAIMER

This chapter is an extended version of the article published by the same author(s) in the following journal.

British Journal of Mathematics & Computer Science, 11(3): 1-8, 2015.

Reviewers' Information

- (1) Anonymous, Universidad Autnoma Metropolitana, Mexico.
- (2) Anonymous, Neijiang Normal University, China.
- (3) Anonymous, Chang-Jung Christian University, Taiwan.